



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Shrikes of Minnesota.—Not exactly knowing the true position held by the Shrikes of Minnesota, I submitted a small series of four to Mr. Ridgway for identification, two from Minneapolis, which were dark colored specimens, and two, which were much lighter, from Lanesboro in the southern part of the State.

The following was the result of the examination. He says: "None of your Shrikes are typical of either *ludovicianus* or *excubitorides*, all being intermediate between the two forms. The specimen I have marked 'A' [one of those from Minneapolis] comes nearest the former, but is not dark enough in the coloration of the upper parts; the others come decidedly nearer *excubitorides* than *ludovicianus*, but are not pale enough to be typical."—GEO. G. CANTWELL, *Lake Mills, Wis.*

Notes on some Minnesota Birds.—It is thought that sufficient interest attaches to the occurrence in Minnesota of the five species of birds mentioned below to justify the publication at this time of the following notes in regard to them. Two of the five—the Burrowing Owl and Henslow's Sparrow—are here reported from the State for the first time, so far as the writer can discover. The published statements in regard to the others have been in such general terms as to give no very definite idea of the real manner of their occurrence.

Colymbus holbœlii. HOLBÆLL'S GREBE.—This bird is found in limited numbers during the summer season in the west-central part of the State. It may, and doubtless does, occur in other portions of the prairie region of the State, but as yet no positive evidence of its presence has been obtained. Birds supposed to be of this species were seen in Elbow Lake, Grant Co., in June, 1879, by Mr. Franklin Benner and the writer, but no positive identification could be then made. Its eggs, taken in the vicinity, were seen at that time in the collection of Mr. Jasper N. Sanford of the town of Elbow Lake. Subsequently, in the fall of 1881, in response to numerous inquiries, Mr. Sanford kindly sent to me the skin of a male of this species together with several eggs taken early in the preceding summer near Elbow Lake. These specimens are now in my collection. This affords satisfactory assurance of the breeding of this Grebe in Minnesota.

Speotyto cunicularia hypogæa. BURROWING OWL.—On July 19, 1881, while passing through the southwestern part of Swift County, which lies in the western part of Minnesota, I came upon a small colony of these Owls living in fox or badger dens in a prairie hillside. There were apparently not more than two pairs of old birds with the young of the year. One of the young birds was shot, and the skin preserved. Many weeks spent in travelling through the prairie portions of the State failed to disclose the presence of this bird in any other locality.

Ammodramus henslowii. HENSLOW'S SPARROW.—A male bird in full song was taken by the writer on June 16, 1880, near Minneapolis, and the species was evidently breeding in the wet marsh where it was shot. In February, 1881, I was shown a specimen by Mr. W. W. Eager which he